

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

CLAIMS

I claim:

1. A method of improving the sexual performance in a male subject, comprising:  
percutaneously delivering a pharmaceutically effective amount of a steroid in the  
5 testosterone synthetic pathway to the subject via a pharmaceutical composition comprising  
the steroid, at least one of a C1-C4 alcohol, and a penetration enhancer.
2. The method in claim 1 wherein the penetration enhancer comprises at least one of  
a C8-C22 fatty acid.
3. The method in claim 1 wherein the fatty acid comprises an alkyl chain length of at  
10 least 12 carbon atoms.
4. The method in claim 1 wherein the lower alcohol comprises at least one of ethanol,  
2-propanol, n-propanol, and mixtures thereof.
5. The method in claim 1 wherein the the steroid is testosterone and the enhancer is  
isopropyl myristate.
- 15 6. The method of claim 5 wherein the composition comprises about 1.0 g w/w of  
testosterone.
7. The method of claim 5 wherein the enhancer comprises about 0.5 g w/w of  
isopropyl myristate.
8. The method of claim 5 wherein the thickener is CARBOPOL.
- 20 9. The method of claim 1 where the steroid comprises about 0.5 g to about 5.0 g  
testosterone, the thickener comprises about 0.10 g to about 2 g of CARBOPOL, the  
enhancer comprises about 0.1 g to about 2 g of isopropyl myristate, the C1-C4 alcohol  
comprises about 40.0 g to about 90 g of ethanol.
10. The method in claim 1 wherein the men are hypogonadal.

11. The method in claim 10 wherein the men suffer from primary hypogonadism.
12. The method in claim 1 wherein the delivering occurs daily.
13. The method in claim 12 wherein the delivery comprises administering the composition to the right/left upper arms/shoulders and to the right/left sides of the abdomen once per day on alternate days.
14. The method in claim 1 wherein the pharmaceutically effective amount of steroid comprises 75 mg of testosterone per day.
15. The method in claim 14 wherein the men achieve hormonal steady state levels of testosterone.
16. The method in claim 1 wherein the improving sexual performance comprises treating impotence in the men.
17. A method of increasing the libido of men comprising:  
delivering a pharmaceutically effective amount of a steroid in the testosterone synthetic pathway to the men percutaneously in a composition comprised of a C1-C4 alcohol, a penetration enhancer, a thickener, testosterone, and water.
18. The method in claim 17 wherein the penetration enhancer comprises at least one of a C8-C22 fatty acid.
19. The method in claim 17 wherein the fatty acid comprises an alkyl chain length of at least 12 carbon atoms.
20. The method in claim 17 wherein the lower alcohol comprises at least one of ethanol, 2-propanol, n-propanol, and mixtures thereof.
21. The method in claim 17 wherein the steroid is testosterone and the enhancer is isopropyl myristate.

22. The method of claim 21 wherein the composition comprises about 1.0 g w/w of testosterone.
23. The method of claim 21 wherein the enhancer comprises about 0.5 g w/w of isopropyl myristate.
- 5 24. The method of claim 21 wherein the thickener is CARBOPOL.
25. The method of claim 17 where the steroid comprises about 0.5 g to about 5.0 g testosterone, the thickener comprises about 0.10 g to about 2 g of CARBOPOL, the enhancer comprises about 0.1 g to about 2 g of isopropyl myristate, the C1-C4 alcohol comprises about 40.0 g to about 90 g of ethanol.
- 10 26. The method in claim 17 wherein the men are hypogonadal.
27. The method in claim 17 wherein the men suffer from primary hypogonadism.
28. The method in claim 17 wherein the delivering occurs daily.
29. The method in claim 28 wherein the delivery comprises administering the composition to the right/left upper arms/shoulders and to the right/left sides of the  
15 abdomen once per day on alternate days.
30. The method in claim 17 wherein the pharmaceutically effective amount of a steroid in the testosterone synthetic pathway comprises 75 mg of testosterone per day.
31. The method in claim 17 wherein the men achieve hormonal steady state levels of testosterone.
- 20 32. A method for improving the efficacy of a pharmaceutical useful for treating erectile dysfunction in a male subject, comprising:
- percutaneously delivering a pharmaceutically effective amount of a steroid in the testosterone synthetic pathway to the subject in a composition comprising at least one of a C1-C4 alcohol, the steroid, and a penetration enhancer; and

administering the pharmaceutical to the subject.

33. The method in claim 32 wherein the subject is eugonadal.
34. The method of claim 32 wherein the pharmaceutical is a phosphodiesterase type 5 inhibitor.
- 5 35. The method in claim 32 wherein the pharmaceutical is at least one of sildenafil citrate, pentoxifylline, yohimbine, apomorphine, alprostadil, papavaerine, phentolamine, and combinations, salts, derivatives and enantiomers of thereof
- 36 36. The method in claim 32 wherein the pharmaceuticals are selected from the group consisting of VIAGRA, UPRIMA, TRENTAL or ACTIBINE.
- 10 37. The method in claim 32 wherein the penetration enhancer comprises at least one of  
37 a C8-C22 fatty acid.
- 38 38. The method in claim 32 wherein the fatty acid comprises an alkyl chain length of at least 12 carbon atoms.
- 39 39. The method in claim 32 wherein the lower alcohol comprises at least one of  
15 ethanol, 2-propanol, n-propanol, and mixtures thereof.
- 40 40. The method in claim 32 wherein the steroid is testosterone and the enhancer is isopropyl myristate.
- 41 41. The method of claim 38 wherein the composition comprises about 1.0 g w/w of testosterone.
- 20 42. The method of claim 38 wherein the enhancer comprises about 0.5 g w/w of  
42 isopropyl myristate.
- 43 43. The method of claim 38 wherein the thickener is CARBOPOL.

- 44 42. The method of claim 32 where the steroid comprises about 0.5 g to about 5.0 g testosterone, the thickener comprises about 0.10 g to about 2 g of CARBOPOL, the enhancer comprises about 0.1 g to about 2 g of isopropyl myristate, the C1-C4 alcohol comprises about 40.0 g to about 90 g of ethanol.
- 5 45 43. The method in claim 32 wherein the men are hypogonadal.
- 46 44. The method in claim 32 wherein the men suffer from primary hypogonadism.
- 47 45. The method in claim 32 wherein the delivering occurs daily.
- 48 46. The method in claim 32 wherein the delivery comprises administering the composition to the right/left upper arms/shoulders and to the right/left sides of the abdomen once per day on alternate days.
- 10 49 47. The method in claim 32 wherein the pharmaceutically effective amount steroid comprises 75 mg of testosterone per day.
- 50 48. The method in claim 32 wherein the men achieve hormonal steady state levels of testosterone.
- 15 51 49. A kit comprised of a pharmaceutical useful for treating erectile dysfunction in a man and a transdermal testosterone gel.
- 52 50. The kit of claim <sup>51</sup>49 wherein the pharmaceutical is a phosphodiesterase type 5 inhibitor.
- 53 51. The kit in claim <sup>51</sup>49 wherein the pharmaceutical is at least one of sildenafil citrate, pentoxifylline, yohimbine, apomorphine, alprostadil, papavaerine, phentolamine, and combinations, salts, derivatives and enantiomers of thereof
- 20 54 52. The kit in claim <sup>51</sup>49 wherein the pharmaceuticals are selected from the group consisting of VIAGRA, UPRIMA, TRENTAL or ACTIBINE.

<sup>53</sup> ~~53~~. The kit of claim <sup>51</sup>~~49~~ wherein the testosterone gel comprises about 0.5 g to about 5.0 g testosterone, a thickener, an enhancer, and a C1-C4 alcohol.

<sup>52</sup> ~~54~~. The kit of claim <sup>55</sup>~~53~~ wherein the thickener comprises about 0.10 g to about 2 g of CARBOPOL, the enhancer comprises about 0.1 g to about 2 g of isopropyl

5 myristate, the C1-C4 alcohol comprises about 40.0 g to about 90 g of ethanol.